



Informed consent form

Op. Dr. Oytun İdil (Estetik plastik cerrahi) / + 90 533 5690649 / oytunmd@gmail.com

PATIENT INFORMATION AND CONSENT FORM **BURIED PENIS SURGERY** **(buried or hidden penis surgery)**

PATIENT'S NAME, SURNAME:

AGE:

DATE OF INFORMATION:

ADDRESS:

This form will inform you about the risks, process and results of IMBURNED PENIS surgery, and will enlighten you about alternative treatments. Please read carefully, sign each page, ask your doctor if you have questions or do not understand the points.

This surgery is performed to remove the penis embedded in the tissue. It is usually performed together with penis enlargement surgery. At the end of the operation, there will be a zigzag or inverted "Y" shaped suture mark on the skin due to the extension on the dorsal part of the base of the penis. If there is a penoscrotal web and it has been opened during the operation, there will be a zigzag stitch mark on the lower part of the penis. These zigzagged scars are due to z-plasties performed to lengthen the skin. The scar that will remain on the lower part of the penis and testicles is not very prominent as a feature of the skin of that area. If there is fat accumulation around the base of the penis, this fat mass is also removed by liposuction (vacuum fat removal).

The only surgical way to remove the implanted penis is to cut the suspensory ligament, lengthen it and remove the excess tissue around the penis. Other non-surgical methods such as tying weights or vacuum pumps are usually not effective.

BEFORE

SURGERY

The anaesthetist will evaluate the patient before surgery. He will examine and ask some questions. He/she will carry out the tests and investigations he/she deems necessary to avoid any problems. Tell your doctor if you have a known or suspected disease (such as heart disease, high blood pressure, diabetes), a symptom or an allergy (such as to

medicines, medical materials such as plasters or food). Describe any previous operations, procedures or medical problems you have had.

ANESTHESIA

This operation is performed under general anaesthesia. In the operating theatre, the patient's heart rate and blood oxygen level are continuously monitored by electronic devices during the operation. Allergy or drug reactions to the materials used are rare and can be fatal. Moreover, people who are sensitive to these substances cannot be identified in advance by routine tests. However, when these undesirable situations occur under hospital conditions, they can be successfully treated and the possibility of harming the patient is extremely small.

OPERATION

PROCESS

Before general anaesthesia, the patient will not eat or drink anything for 6-8 hours with an empty stomach. After the operation, nothing will be given by mouth for at least 4-6 hours. These periods can be changed by the patient's doctor. The duration of the operation lasts 1.5 - 4 hours depending on the procedure to be performed. After the operation, the patient is usually monitored in the waking room for 1 hour and then sent to his/her room. However, if nausea and similar problems occur due to anaesthesia, the duration of stay may be prolonged.

POSSIBLE RISKS ASSOCIATED WITH SURGERY

As after any surgery, there may be some undesirable conditions (complications) after this operation. Infection, bleeding, asymmetry, melting of the injected fat tissue, loss of stabilisation in the penis (more movement of the penis from left to right), loss of sensation, necrosis may occur.

In order to reduce the likelihood of problems, antibiotics, painkillers and other medications recommended by the doctor should be used without interruption after the operation, and the dressing or bandage should be applied regularly. In case of any problem related to the surgery, the doctor who performed the surgery should be consulted first, since he/she is the person who can best evaluate the source and solution of the problem.

Smoking and tobacco use adversely affect tissue circulation and significantly increase the possibility of all problems that may occur.

RISKS OF LIPOSACCHIN (VACUUM FAT REMOVAL) AND FAT INJECTION (in buried penile deformity)

Patients with loose skin, reduced skin tension, other medical problems (such as diabetes, heart disease, collagen tissue disease, vascular disease), unrealistic desires (such as a very fat patient expecting ideal body measurements) are not suitable for this surgery. Postoperative infection (inflammation) may occur. Most of the time it improves with dressing and medication. However, this may lead to skin shrinkage and contour disorders.

Necessary precautions are taken against the risk of infection. There may also be haematoma (blood collection) in the operation areas. In order to prevent bleeding during the operation, a special bleeding-reducing solution is injected into all liposuction areas and then the procedure begins. If a haematoma is formed, subsequent intervention (such as insertion of tubes called resistors through the cannula holes) may be required to drain it. If bleeding is excessive, it may be necessary to give blood. Swelling and bruising are expected in the relevant areas after the operation. These may last for several weeks. Seroma (serum collection) may also occur at the operation site. There is no risk of this. It is usually absorbed back by the body after a while.

Although it is a very weak possibility, an allergic reaction to the drugs and suture materials used may develop. In this surgery, sensory disturbances, fluctuations, collapses, heights, symmetry disorders between the two sides may occur. These may be temporary or permanent. There is a risk of malfunction of another organ or vessel (embolism) as a result of fat escaping into a large vessel opened in liposuction (vacuum fat removal) surgery. When this embolism travels to the brain, heart and lungs, it can lead to serious conditions that can lead to death. This risk is minimised by administering an anti-bleeding and vasoconstrictive solution (tumescent technique) to the operation area before the procedure. Skin loss is another very rare complication of liposuction. The fat melts a little in the area where it is injected. The fat injection may need to be repeated. As far as it is known, melting can be up to 40% of the injected volume. In order to reduce the possibility of problems, antibiotics, painkillers and other medications recommended by the doctor should be used without interruption, dressings, bandages and special corsets should be applied regularly. In case of any problem related to the surgery, the doctor who performed the surgery should be consulted first, since he/she is the person who can best evaluate the source and solution of the problem. Smoking and tobacco use adversely affect tissue circulation and significantly increase the possibility of all problems that may occur.

PATIENT

CONSENT

My doctor told me about my problems, the type of treatment to be performed, my surgery, alternative treatments, risks and complications I might encounter. He gave me the opportunity to ask any questions I had. I understand that I can be given serum and, if necessary, blood during the operation. I know that I can be operated again under anaesthesia if complications that may require surgical procedure develop. I know that if surgical procedure is required due to complications, I know that I will have to pay the hospital costs of these additional surgical

procedures.

I understand that tissues and organs may be removed from me during the operation to remove excess skin and soft tissues, to be used in another area (called graft or flap) or for pathological examination.

I understand that scars will remain after the operation and that they will not disappear for life. I agree to do what is recommended (massage with special creams, application of silicone sheets) to keep these scars as minimal and as good as possible. I authorise the taking of medical pictures and videos of me before, during and after the operation. I understand that these pictures and videos may be shown to other doctors for medical and scientific purposes. I understand that no payment will be made to me for the use of these images and videos in medical publications (articles or books). I understand that there are also risks associated with general anaesthesia, that these are the responsibility of the anaesthesiologist, and that I will be explained separately and a form will be signed.

As a result, I understand that I am not guaranteed the outcome of this operation, that there is a possibility that the expected result may not be achieved for various reasons, and that even worse results may be obtained than the current situation. This consent form may contain errors or omissions. If I have any questions about this form or the surgery, I know that I can consult my doctor via my doctor's e-mail address (oytunmd@gmail.com), mobile phone (+90 533 5690649 and +90 505 2965569) and whatsapp (+90 533 5690649).

**Information about the Covid-19 pandemic:
I have been informed that the surgeries were performed during the covid-19 pandemic but by taking all necessary precautions. Despite all these precautions, knowing that I can get covid-19 infection outside the hospital, I agree to be operated by taking the risks into consideration.**

At the end of these, I accept the application of the "buried penis" surgery explained to me in detail by my doctor Op. Dr. Oytun idil with my own consent and without any influence.

Date: /..... / **2023**

Patient's **name** **and** **surname:**

Signature: (please sign by writing "I READ, I UNDERSTOOD")

Relative/witness, **name** **and** **surname:**

The patient has signed the form in my presence, awake and conscious.

Signature: